



ISSN 1005-264X
www.plant-ecology.com

植物生态学报

Chinese Journal of Plant Ecology

第43卷 第6期 2019年6月 Vol. 43 No. 6 June 2019



主办单位：中国科学院植物研究所
中国植物学会

Sponsors: Institute of Botany, Chinese Academy of Sciences
Botanical Society of China

植物生态学报

Zhiwu Shengtai Xuebao

2019年6月 第43卷 第6期

目次

综述	赵丹丹 马红媛 李阳 魏继平 王志春
471 磷影响陆地生态系统碳循环过程及模型表达方法 黄玫 王娜 王昭生 巩贺	512 泥炭地特征性环境因子促进泥炭藓持久孢子库的形成 冯璐 卜兆君 吴玉环 刘莎莎 刘超
研究论文	521 秦岭东段栓皮栎枝条非结构性碳水化合物含量的季节动态 章异平 海旭莹 徐军亮 吴文霞 曹鹏鹤 安文静
482 神农架常绿落叶阔叶混交林碳氮磷化学计量比 刘璐 葛结林 舒化伟 赵常明 徐文婷 申国珍 谢宗强	532 全球森林土壤微生物生物量碳氮磷化学计量的季节动态 李品 木勒德尔·吐尔汗拜 田地 冯兆忠
490 2000–2017年新疆天山植被水分利用效率时空特征及其与气候因子关系分析 艾则孜提约麦尔·麦麦提 玉素甫江·如素力 何辉 拜合提尼沙·阿不都克日木	543 入侵植物加拿大一枝黄花根际解钾菌多样性及解钾活性 闫雅楠 叶小齐 吴明 闫明 张昕丽
501 水分和养分添加对羊草功能性状和地上生物量的影响	

封面说明: 新疆巴音郭楞蒙古自治州的巴音布鲁克草原景观(新疆师范大学地理科学与旅游学院孙浩捷摄)。巴音布鲁克草原位于天山山脉中段的高山间盆地,四周为雪山环抱的典型禾草草甸草原。艾则孜提约麦尔·麦麦提等基于MODIS数据,研究了新疆天山及其南北两侧主要绿洲的植被水分利用效率(WUE)及其时空变化,探讨了其影响因素(本期490–500页)。

Chinese Journal of Plant Ecology

June 2019 Vol. 43 No. 6

CONTENTS

Review

- 471 Modeling phosphorus effects on the carbon cycle in terrestrial ecosystems
HUANG Mei, WANG Na, WANG Zhao-Sheng, and GONG He

Research Articles

- 482 C, N and P stoichiometric ratios in mixed evergreen and deciduous broadleaved forests in Shennongjia, China
LIU Lu, GE Jie-Lin, SHU Hua-Wei, ZHAO Chang-Ming, XU Wen-Ting, SHEN Guo-Zhen, and XIE Zong-Qiang
- 490 Spatio-temporal characteristics of vegetation water use efficiency and its relationship with climate factors in Tianshan Mountains in Xinjiang from 2000 to 2017
Aizezitiyuemaier MAIMAITI, Yusufujiang RUSULI, HE Hui, and Baihetinisha ABUDUKERIMU
- 501 Effects of water and nutrient additions on functional traits and aboveground biomass of *Leymus chinensis*

ZHAO Dan-Dan, MA Hong-Yuan, LI Yang, WEI Ji-Ping, and WANG Zhi-Chun

- 512 Characteristic environmental factors in peatlands facilitate the formation of persistent *Sphagnum* spore banks

FENG Lu, BU Zhao-Jun, WU Yu-Huan, LIU Sha-Sha, and LIU Chao

- 521 Seasonal dynamics of non-structural carbohydrate content in branch of *Quercus variabilis* growing in east Qinling Mountain range

ZHANG Yi-Ping, HAI Xu-Ying, XU Jun-Liang, WU Wen-Xia, CAO Peng-He, and AN Wen-Jing

- 532 Seasonal dynamics of soil microbial biomass carbon, nitrogen and phosphorus stoichiometry across global forest ecosystems

LI Pin, Muledeer TUERHANBAI, TIAN Di, and FENG Zhao-Zhong

- 543 Diversity and potassium-solubilizing activity of rhizosphere potassium-solubilizing bacteria of invasive *Solidago canadensis*

YAN Ya-Nan, YE Xiao-Qi, WU Ming, YAN Ming, and ZHANG Xin-Li

Cover illustration: The Bayanbulak grassland landscape in the Bayingolin Mongol Autonomous Prefecture, Xinjiang, China (Photographed by SUN Hao-Jie, Institute of Geographical Science and Tourism, Xinjiang Normal University). The Bayanbulak grassland is located in intermountain basin covered by meadow grasses. Maimaiti *et al.* studied the spatio-temporal distribution of the vegetation water use efficiencies (*WUE*) in Tianshan Mountains in Xinjiang and its north and south sites and their influencing factors (Pages 490–500 of this issue).