


白天道, 余春兰, 甘泽朝, 赖海荣, 杨隐超, 黄厚宸, 蒋维昕 (2020). 细叶云南松种实性状变异与地理气象因子的关联. 植物生态学报, 44, 1224–1235. DOI: 10.17521/cjpe.2020.0269

Bai TD, Yu CL, Gan ZC, Lai HR, Yang YC, Huang HC, Jiang WX (2020). Association of cone and seed traits of *Pinus yunnanensis* var. *tenuifolia* with geo-meteorological factors. *Chinese Journal of Plant Ecology*, 44, 1224–1235. DOI: 10.17521/cjpe.2020.0269

<https://www.plant-ecology.com/CN/10.17521/cjpe.2020.0269>

附录I 细叶云南松天然种群采样点(▲)及用于空间插值分析的气象站点(●)分布图

Supplement I Map of sampling sites (▲) and the weather stations (●) for spatial interpolation of eight natural populations in *Pinus yunnanensis* var. *tenuifolia*



BW, 兴义坝汪; BY, 册亨弼佑; CJ, 兴义岔江; DG, 望谟大观; DT, 罗甸大亭; QX, 安龙钱相; WJ, 罗甸伍家坟; XQ, 安龙新桥。曲线代表区域内分布的二级(蓝色)及三级(红色)河流。

BW, Bawang, Xingyi; BY, Biyou, Ceheng; CJ, Chajiang, Xingyi; DG, Dagan, Wangmo; DT, Dating, Luodian; QX, Qianxiang, Anlong; WJ, Wujiafen, Luodian; XQ, Xinqiao, Anlong. The blue and red curves represent the rivers in the distribution area.